


#### Ice Climatology & Modeling Team

#### Team Members:

Anne Clites (GLERL), Xuezhi Bai (CILER), Haoguo Hu (CILER), Ray Assel, (CILER), Yi Yang (CILER)

Ayumi Fujisaki, (CILER), Lin Luo (CILER)



- 1) Investigates relationships between lake ice / regional climate variability and changing climate patterns.
- 2) Develops and implements coupled ice-circulation-ecosystem models to simulate ice-circulation-ecosystem's response to a changing climate, and to develop a capability for downscaling and projecting climate.
  scenarios into Great Lakes regional scales.
- 3) Conducts interdisciplinary research and tests hypotheses using models along with in situ and satellite measurements.

2

# **Key Scientific Questions**



3

- 1) Which large-scale climate patterns (forcings) drive seasonal to decadal variability of lake ice and regional climate?
- 2) What factors determine lake ice predictability on time scales from season to decades using statistical methods?
- 3) How does shrinking lake ice relate to lake level variability and lake water temperature?
- 4) Can we predict lake ice on synoptic time scales using a coupled Great Lakes Ice-circulation Model (GLIM)?



Which climate patterns (forcings) drive seasonal to decadal variability of lake ice & regional climate?

North Atlantic Oscillation (NAO)

(Arctic Oscillation)


Pacific North America Pattern
(El Nino/La Nina, ENSO)

Linear relationship

Ice and ENSO:
Nonlinear relationship

The Great Lakes are not the action center – we are on the periphery of the action center.

Bai et al. (submitted to JGR) Wang et al. EOS, 2010



These are two examples of graphs of Lakes Erie and Ontario with similar size surface areas but different depths.

For deep lakes standard deviation is equal to or greater than the mean, which means it is poor predictor of long-term ice.

Wang et al. submitted to J. Climate

Bai et al. submitted to JGR


What factors determine lake ice predictability from season to decades using statistical methods? For non-linear regression model: Y=0.3-0.118xNino3.4-0.299xNino3.42 -0.208xNAO Coverage Observation r(model\_nino3.4\*\*2,obs.)=0.5 2 r(moce/\_nino3.4,obs.)=0.39 1.5 0.5 Normalized Ice Nonlinear -1.5 -2-2.5 -31965 1970 1975 1980 1985 1990 1995 2000 2005 year

These are two examples of graphs of Lake Erie and Lake Ontario with similar size surface areas but different depths.

For deep lakes standard deviation is equal to or greater than the mean, which means it is poor predictor of long-term ice.

Wang et al. submitted to J. Climate

Bai et al. submitted to JGR




#### Hypothesis:

Water level ~


- + a Ice (-Ta/Tw)
- b E(Ta/Tw, -lce)
- + c P
- + d Runoff (land process~Ta/Tl, Veg., land-air energy budget)

Need a coupled model to test it! Wang et al. submitted to J. Climate



Test in Lake Superior


Wang et al. GRL, 2009



Without lake ice in a model, the simulated water level using water only model is over-shot or overestimated; similarly, without sea ice in a model, the simulated lake water temperature using a water only model is over-heated.


Wang et al. 2010; JGLR

9



GLIM produces reasonable lake ice cover in comparison to satellite measurement.

Wang et al. 2010, JGLR




Overall comparison between the simulated ice area and measured ice area by satellite is very good. (Wang et al. 2010, JGLR).

This is one of the NOAA's Climate Goals: Ice forecasting.

Potential users: Industries, U.S. Coast Guard, public/private, governments, academia. Collaborators: National Weather Service, National Ice Center, University of Michigan.

11



http://www.glerl.noaa.gov/data/ice/atlas/

## **Delivered Tools, Products and Services**


- Great Lakes Ice-circulation Model (GLIM) delivered to GLCFS, (2007-present)
- Unstructured-grid, entire Great Lakes watershed-scale GLIM for climate studies: Hindcast and downscaling simulations, projection of climate scenarios into GL watershed scale

(to be coupled to GL WRF (regional climate model), ecosystems model, and other subsystem models to develop capability for climate studies in line with National Climate Service)



http://www.glerl.noaa.gov/data/ice/atlas/


13



The update of the Great Lakes Ice Atlases are necessary and advance our understanding of climate change (Wang et al. 2010, submitted to J. Climate).

This is one of the NOAA's Climate Goals: Ice forecast.

Potential users: Industries, U.S. Coast Guard, public/private, governments, academia.



High-resolution, coupled GLIM on a Great Lakes watershed scale is essential for regional climate simulation, downscaling, and projection.


15

### **Future Directions**



- Continue interdisciplinary research associated with lake ice and climate
- Continue to improve Great Lakes Ice-circulation Model (GLIM) for Great Lakes Coastal Forecasting System
- Upgrade Great Lakes Ice Atlas to web-based, user-friendly Great Lakes Ice Mapping System
- Develop unstructured-grid integrated ice-circulationecosystem modeling capability for climate studies
- Couple to atmosphere and hydrology to develop regional Great Lakes Earth System Model





### Further information:

<u>Jia.wang@noaa.gov</u> http://www.glerl.noaa.gov/about/pers/profiles/wang.html